



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## BiFORCE 100SC Termiticide & Insecticide

ACTIVE CONSTITUENT: 100g/L BIFENTHRIN

**GROUP 3A INSECTICIDE**

For the pre- and post- construction management of subterranean termites in accord with AS 3660 and for the control of termites and a range of interior and exterior urban pests as specified in the Directions For Use Table.

**IMPORTANT: RESTRICTED CHEMICAL PRODUCT ONLY TO BE SUPPLIED TO, OR USED BY AN AUTHORISED PERSON**

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING**



Distributed by:

**Sherwood Chemicals**  
25 Valley Road  
Wembley Downs WA 6019  
Tel: 0421 667 972

**CONTENTS 1L APVMA Approval No: 62837/1L/0109**

**BiFORCE 100SC Termiticide & Insecticide** is a Water-based Suspension Concentrate formulation

### CONDITIONS OF USE BY AUTHORISED PERSONS

The Pest Control Operator must be licensed under state legislation.

The Pest Control Operator must notify site supervisor, if any, and workers who come into contact with uncovered treated soil prior to laying the moisture membrane, to wear appropriate personal protective equipment and to observe re-entry requirements.

(For personal protective equipment, refer to "SAFETY DIRECTIONS", and for re-entry refer to "PRECAUTIONS AND RE-ENTRY PERIOD" below).

### PRECAUTIONS AND RE-ENTRY PERIOD

DO NOT spray directly on humans, pets, or animals. Avoid contact with food, food utensils or preparation surfaces.

#### Re-entry period

**Pre-construction: Re-entry** : - DO NOT allow entry into uncovered treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after use daily.

**Post-construction and urban pest control:** Allow treated areas to completely dry (normally 3-4 hours) and ventilate buildings before re-occupying. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after use daily.

### STORAGE, SPILLAGE AND DISPOSAL

Store in closed original container, in a cool, well-ventilated area away from children, animals, food and feedstuffs. DO NOT store for prolonged periods in direct sunlight.

In case of spillage, confine and absorb spilled product with absorbent material such as sand, clay or cat litter. Dispose of waste as indicated below or according to the Australian Standard AS 2507 - Storage and Handling of Pesticides.

DO NOT allow spilled product to enter sewers, drains, creeks or any other waterways.

#### For REFILLABLE containers

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### For 1L Containers

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt. DO NOT bury waste or surplus product. Dispose of undiluted waste by either dilution and use according to the Directions for Use or returning to the point of purchase in the original container for controlled disposal. Dispose of diluted surplus product by using according to the Directions for Use. DO NOT re-use empty container.

### SAFETY DIRECTIONS

#### Poisonous if swallowed.

May irritate eyes and skin. Repeated exposures may cause allergic disorders. Avoid contact with eyes and skin.

#### For termite control in buildings

**and structures:** When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PCV, or nitrile gloves. After each days use, wash gloves and contaminated clothing.

**For handheld application:** When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC or nitrile gloves.

When using the prepared spray wear protective waterproof clothing, elbow length PVC or nitrile gloves and water resistant footwear. After each days use, wash gloves and contaminated clothing. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.


#### MATERIAL SAFETY DATA SHEET.

Additional information is listed in the Material Safety Data Sheet (MSDS). Contact Sherwood Chemicals for a copy of the MSDS.

Tel: 0421 667 972.

### CONDITIONS OF SALE:

The use of BiFORCE 100SC Termiticide & Insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by Sherwood Chemicals regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Sherwood Chemicals accepts no responsibility for any consequence whatsoever resulting from the use of this product.



BiFORCE 100SC  
Batch No:  
DOM:

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# BiFORCE 100SC

## Termiticide & Insecticide



ACTIVE CONSTITUENT: 100g/L BIFENTHRIN

GROUP **3A** INSECTICIDE

For the pre- and post- construction management of subterranean termites in accord with AS 3660 and for the control of termites and a range of interior and exterior urban pests as specified in the Directions For Use Table.

**IMPORTANT:**  
RESTRICTED CHEMICAL PRODUCT ONLY TO BE SUPPLIED  
TO, OR USED BY AN AUTHORISED PERSON  
**IMPORTANT:**  
READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



Distributed by:  
**Sherwood Chemicals**  
25 Valley Road  
Wembley Downs WA 6019  
Tel: 0421 667 972



CONTENTS 5L  
APVMA Approval No: 62837/5L/0109

File Name: 62837\_44086\_5L\_LABEL\_FRONT\_MPL\_V4 Dimensions:105.00mm (W) x 155.00mm (H) Scale of Print: A4 100% A3 100%

**BiFORCE 100SC Termiticide & Insecticide** is a Water-based Suspension Concentrate formulation

#### CONDITIONS OF USE BY AUTHORISED PERSONS

The Pest Control Operator must be licensed under state legislation. The Pest Control Operator must notify site supervisor, if any, and workers who come into contact with uncovered treated soil prior to laying the moisture membrane, to wear appropriate personal protective equipment and to observe re-entry requirements. (For personal protective equipment, refer to "SAFETY DIRECTIONS", and for re-entry refer to "PRECAUTIONS AND RE-ENTRY PERIOD" below).

#### PRECAUTIONS AND RE-ENTRY PERIOD

DO NOT spray directly on humans, pets, or animals. Avoid contact with food, food utensils or preparation surfaces.

#### Re-entry period

**Pre-construction: Re-entry:** - DO NOT allow entry into uncovered treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after use daily.

**Post-construction and urban pest control:** Allow treated areas to completely dry (normally 3-4 hours) and ventilate buildings before re-occupying. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after use daily.

#### STORAGE, SPILLAGE AND DISPOSAL

Store in closed original container, in a cool, well-ventilated area away from children, animals, food and feed stuffs.

DO NOT store for prolonged periods in direct sunlight.

In case of spillage, confine and absorb spilled product with absorbent material such as sand, clay or cat litter. Dispose of waste as indicated below or according to the Australian Standard AS 2507 - Storage and Handling of Pesticides.

DO NOT allow spilled product to enter sewers, drains, creeks or any other waterways.

#### Disposal for 5L Containers

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots.

Empty containers and product should not be burnt. DO NOT bury waste or surplus product.

Dispose of undiluted waste by either dilution and use according to the Directions for Use or returning to the point of purchase in the original container for controlled disposal. Dispose of diluted surplus product by using according to the Directions for Use. DO NOT re-use empty container.

#### SAFETY DIRECTIONS

**Poisonous if swallowed.** May irritate eyes and skin. Repeated exposures may cause allergic disorders. Avoid contact with eyes and skin.

**For termite control in buildings and structures:** When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC, or nitrile gloves. After each days use, wash gloves and contaminated clothing.

**For handheld application:** When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC or nitrile gloves.

When using the prepared spray wear protective waterproof clothing, elbow length PVC or nitrile gloves and water resistant footwear. After each days use, wash gloves and contaminated clothing. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

#### MATERIAL SAFETY DATA SHEET.

Additional information is listed in the Material Safety Data Sheet (MSDS). Contact Sherwood Chemicals for a copy of the MSDS. Tel: 0421 667 972.

#### CONDITIONS OF SALE:

The use of BiFORCE 100SC Termiticide & Insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by Sherwood Chemicals, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Sherwood Chemicals accepts no responsibility for any consequence whatsoever resulting from the use of this product.

CONTENTS 5L

Batch No:

DOM:

File Name: 62837\_44086\_5L\_LABEL\_BACK\_MPL\_V4 Dimensions:105.00mm (W) x 155.00mm (H) Scale of Print: A4 100% A3 100%

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# BiFORCE 100SC

## Termiticide & Insecticide



ACTIVE CONSTITUENT: 100g/L BIFENTHRIN

**GROUP 3A INSECTICIDE**

For the pre- and post- construction management of subterranean termites in accord with AS 3660 and for the control of termites and a range of interior and exterior urban pests as specified in the Directions For Use Table.

**IMPORTANT:**  
RESTRICTED CHEMICAL PRODUCT ONLY TO BE SUPPLIED TO, OR USED BY AN AUTHORISED PERSON

**IMPORTANT:**  
READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



Distributed by:  
**Sherwood Chemicals**  
25 Valley Road  
Wembley Downs WA 6019  
Tel: 0421 667 972

- CONTENTS 10L  
APVMA Approval No: 62837/10L/0109
- CONTENTS 20L  
APVMA Approval No: 62837/20L/0109
- CONTENTS 200L  
APVMA Approval No: 62837/200L/0109



**BiFORCE 100SC Termiticide & Insecticide** is a Water-based Suspension Concentrate formulation

#### CONDITIONS OF USE BY AUTHORISED PERSONS

The Pest Control Operator must be licensed under state legislation. The Pest Control Operator must notify site supervisor, if any, and workers who come into contact with uncovered treated soil prior to laying the moisture membrane, to wear appropriate personal protective equipment and to observe re-entry requirements. (For personal protective equipment, refer to "SAFETY DIRECTIONS", and for re-entry refer to "PRECAUTIONS AND RE-ENTRY PERIODS" below).

#### PRECAUTIONS AND RE-ENTRY PERIOD

DO NOT spray directly on humans, pets, or animals. Avoid contact with food, food utensils or preparation surfaces.  
**Re-entry period**

**Pre-construction: Re-entry** : - DO NOT allow entry into uncovered treated areas until the spray has dried.

When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after use daily.

**Post-construction and urban pest control:** Allow treated areas to completely dry (normally 3-4 hours) and ventilate buildings before re-occupying. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear.

#### STORAGE, SPILLAGE AND DISPOSAL

Store in closed original container, in a cool, well-ventilated area away from children, animals, food and feedstuffs. DO NOT store for prolonged periods in direct sunlight.

In case of spillage, confine and absorb spilled product with absorbent material such as sand, clay or cat litter. Dispose of waste as indicated below or according to the Australian Standard AS 2507 - Storage and Handling of Pesticides. DO NOT allow spilled product to enter sewers, drains, creeks or any other waterways.

#### Disposal for 10L, 20L & 200L Containers

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

DO NOT bury waste or surplus product.

Dispose of undiluted waste by either dilution and use according to the Directions for Use or returning to the point of purchase in the original container for controlled disposal. Dispose of diluted surplus product by using according to the Directions for Use.

DO NOT re-use empty container.

#### SAFETY DIRECTIONS

**Poisonous if swallowed.** May irritate eyes and skin. Repeated exposures may cause allergic disorders. Avoid contact with eyes and skin.

**For termite control in buildings and structures:** When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PCV, or nitrile gloves. After each days use, wash gloves and contaminated clothing.

**For handheld application:** When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC or nitrile gloves. When using the prepared spray wear protective waterproof clothing, elbow length PVC or nitrile gloves and water resistant footwear. After each days use, wash gloves and contaminated clothing. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

#### MATERIAL SAFETY DATA SHEET.

Additional information is listed in the Material Safety Data Sheet (MSDS). Contact Sherwood Chemicals for a copy of the MSDS. Tel: 0421 667 972.

#### CONDITIONS OF SALE:

The use of BiFORCE 100SC Termiticide & Insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by Sherwood Chemicals, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Sherwood Chemicals accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Batch No:

DOM:



LEAFLET

## POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# BiFORCE 100SC

## Termiticide & Insecticide

ACTIVE CONSTITUENT: 100g/L BIFENTHRIN

**GROUP 3A INSECTICIDE**

For the pre- and post- construction management of subterranean termites in accord with AS 3660 and for the control of termites and a range of interior and exterior urban pests as specified in the Directions For Use Table.

**IMPORTANT:**  
**RESTRICTED CHEMICAL PRODUCT ONLY TO BE SUPPLIED TO, OR USED BY AN AUTHORISED PERSON**

**IMPORTANT:**  
**READ THIS LEAFLET BEFORE OPENING OR USING**



Distributed by:  
**Sherwood Chemicals**  
25 Valley Road  
Wembley Downs WA 6019  
Tel: 0421 667 972

APVMA Approval Number: 62837/0109

#### Horizontal Barrier Treatments:

Use 5L of solution per m<sup>2</sup> of soil. Apply the termiticide solution evenly to the soil surface area to ensure the provision of a continuous barrier with no gaps. To minimise drift, use low pressure, high volume spray equipment delivering large coarse droplets. On impervious soils where the application of 5L/m<sup>2</sup> would cause excessive run-off, the application volume may be reduced provided the concentration of the solution is increased by a corresponding amount. For example, the volume of applied concentrate must remain constant at 25, 50 or 75mL/m<sup>2</sup> depending on the location and the situation. DO NOT apply solution volumes below 2L/m<sup>2</sup>.

In situations where the soil surface is very dry and conditions are conducive to rapid drying, the areas to be treated should be moistened prior to the termiticide application. It is important to note that when applying a horizontal barrier to the perimeter of a building or structure the chemical barrier is deemed to have a depth of 80mm. In situations where the solution will not readily wet the soil to the required depth, loosen soil to a depth of 80mm by 150mm wide and apply 1.5L of solution per lineal metre.

#### Vertical Barrier Treatments:

To install vertical barrier use a minimum of 100L of solution per m<sup>3</sup> of soil. Vertical barriers must be a minimum of 150mm wide, extend down to 80mm below the top of the footing and be complete and continuous. Vertical barriers can be installed by trenching and treating the soil as it is backfilled, by soil rodding or by the use of certified reticulation systems, as described in the Australian Standard AS 3660 Series. The preferred method of installing a vertical barrier treatment is either by trenching and treating the soil as it is backfilled or by delivery via a certified reticulation system. When using the soil rodding method to establish a vertical barrier the distance between rod spacing should be as per the following table. To improve soil penetration, the soil should be loosened to depth of 150mm.

Soil type	Rod Spacing (mm)
Heavy clay	150
Clay loams	200
Loams	250
Sands	300

#### Perimeter Barrier Treatments:

Perimeter barrier consists of horizontal barriers at least 150mm wide adjoining a vertical barrier of at least 150mm in width. A perimeter barrier must completely surround all buildings, pipes, piers and service penetrations. In buildings with suspended floors with greater than 400mm crawl space, perimeter barriers should be installed to surround piers, stumps and service penetrations and completely about all substructure walls. To ensure provision of a continuous barrier use a minimum of 100L of solution per m<sup>3</sup> of soil. This equates to a delivery volume of 5L of solution per linear metre for a 300mm vertical barrier, or 10L of solution per linear meter for a 600mm vertical barrier. Termites may gain access behind engaged piers against single brick walls unless the soil is treated on both sides of the wall down to the footing.

#### Post-Construction under Slab Treatments:

For concrete slabs, the solution needs to be injected through pre-drilled holes through the slab, at intervals between 150mm and 300mm. The following table shows the recommended hole spacing and recommended volume of spray solution required per hole, depending on the soil type.

Soil type	Hole Spacing (mm)	Litres per hole
Heavy clay	150	1.5
Clay loams	200	2
Loams	250	2.5
Sands	300	3

Application equipment used to inject BiFORCE 100SC Termiticide & Insecticide through pre-drilled holes in an interior situation must be in good working order, free of any leaks and the injector must have the tip shut-off to prevent nozzle dripping. Lateral dispersion tips are recommended. Drill holes must be resealed following injection of the BiFORCE 100SC solution. The decision and/or need for drilling concrete floor slabs should only be made after a thorough inspection of the building. The degree of termite activity should also be taken into consideration.

#### Treatment in Conjunction with Physical Barriers:

In situations where the termite protection system is to consist of a combination of both physical and chemical barriers, each certified system must be installed according to the relevant and appropriate product specification and the Australian Standard AS 3660 Series.

#### CONDITIONS OF SALE

The use of BiFORCE 100SC Termiticide & Insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by Sherwood Chemicals regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, Sherwood Chemicals accept no responsibility for any consequence whatsoever resulting from the use of this product

#### CONDITION OF USE BY AUTHORISED PERSONS

The pest control operator must be licensed under state legislation. The pest operator must notify the site supervisor, if any, and workers who come into contact with uncovered treated soil prior to laying the moisture membrane, to wear appropriate personal protective equipment and to observe re-entry requirements. (For personal protective equipment, refer to "SAFETY DIRECTIONS", and for re-entry, refer to "PRECAUTION: RE-ENTRY PERIODS", below \*\*\*).

#### GENERAL INSTRUCTIONS

**General Pest Control** - BiFORCE 100SC Termiticide & Insecticide is a powerful knockdown and residual pesticide. Ants, cockroaches, fleas, flies, mosquitoes, spiders, ticks and wasps are controlled by direct contact with spray and also by residual action as they come into contact with treated surfaces.

**Termites** - The use of BiFORCE 100SC Termiticide & Insecticide will help prevent and control subterranean termite infestations in and around building and structures when used in accordance with the Australian Standard AS 3660 Series, Termite Management. A dilute termiticidal solution must be adequately dispersed into the soil to establish a barrier between the building and subterranean termites in the soil. The purpose of a termite barrier is to prevent concealed termite entry into the building. The biology and behaviour of the termite species involved should be considered by the pest control operator in determining which control measures are most appropriate to control and prevent termite infestations.

#### INSECTICIDE RESISTANCE WARNING

**GROUP 3A INSECTICIDE**

For insecticide resistance management BiFORCE 100SC is a Group 3A insecticide.

Some naturally occurring insect biotypes resistant to BiFORCE 100SC and other group 3A insecticide may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if BiFORCE 100SC or other group 3A insecticides are used repeatedly. The effectiveness of BiFORCE 100SC on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Sherwood Chemicals accepts no liability for any losses that may result from the failure of BiFORCE 100SC to control resistant insects. BiFORCE 100SC may be subject to specific resistance management strategies. For further information, contact your local supplier or SHERWOOD CHEMICALS representatives or local agricultural department agronomist.

#### MIXING

Add the required quantity BiFORCE 100SC Termiticide & Insecticide to water in the spray tank and mix thoroughly. Maintain agitation during both mixing and application.

To facilitate even application of the termiticide solution over the area to be treated, the addition of a marker dye at label rate is recommended. On hard to wet soil the penetration of the termiticide solution may be improved by the addition of a soil surfactant at label rate.

#### CRITICAL APPLICATION DETAILS

The application of BiFORCE 100SC Termiticide & Insecticide to form both horizontal and vertical chemical barriers must be in accordance with the Australian Standard AS 3660 Series. For treatment of new and existing buildings, both horizontal and vertical barriers may be required around and under the building. External perimeter barriers and where required, internal perimeter barriers, are an essential part of this treatment. The purpose of a chemical termite soil barrier is to provide a continuous, no gap barrier between the building and the termite colony. It is therefore essential that the pest control operator is familiar with the construction details of the building. For further details, refer to the "Horizontal Barrier Treatments" and "Vertical Barrier Treatments" statements in this leaflet and to the Australian Standard AS 3660 Series.

#### SAFETY DIRECTIONS

**Poisonous if swallowed.** May irritate eyes and skin. Repeated exposures may cause allergic disorders. Avoid contact with eyes and skin.

**For termite control in buildings and structures:** When opening the container, preparing spray and using prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length PVC or nitrile gloves. When using the prepared spray wear protective waterproof clothing, elbow length PVC or nitrile gloves and water resistant footwear. After each days use, wash gloves and contaminated clothing. Wash hands after use.

**For handheld application:** When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC or nitrile gloves. When using the prepared spray wear protective waterproof clothing, elbow length PVC or nitrile gloves and water resistant footwear. After each days use, wash gloves and contaminated clothing. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS). A MSDS for BiFORCE 100SC is available from Sherwood Chemicals on request. Call customer service on Tel: 0421 667 972

#### Reticulation systems:

BiFORCE 100SC Termiticide & Insecticide can be used through reticulation systems to form horizontal and vertical barriers under and around structures and all service penetrations. The reticulation system must be certified and be capable of distributing the termiticide solution according to the product label and Australian Standard AS 3660 Series. In situations using reticulations system to form barriers around perimeter and/or service penetrations only, a full pre-construction soil applied BiFORCE 100SC Termiticide & Insecticide horizontal barrier is recommended. It is the responsibility of the builder and all relevant sub-contractors to ensure that all termite barrier systems are installed in accordance with the relevant product installation directions and the Australian Standard AS 3660 Series.

#### Service Requirements:

Service requirements are to be determined as a result of least an annual inspection by a licensed Pest Control Operator. More frequent inspections may be required in high risk termite areas. In determining the need for service, factors such as local termite pressure, breaches of the barrier and termiticide longevity should be considered.

**Subterranean termites are on occasions capable of bridging termite barriers and therefore regular inspections, as detailed in the Australian Standard AS 4349.3 will significantly increase the probability of detection of termite activity before any damage or costly repairs are required.**

Several factors contribute to the longevity of the termite treatment and must be considered when evaluating the need for re-treatment. The actual protection period will depend on the termite hazard, climate, soil conditions and rate of termiticide used. Refer to Table B for the expected protection periods provided.

#### \*\*\*PRECAUTIONS AND RE-ENTRY PERIOD

DO NOT spray directly on humans, pets or animals. Avoid contact with food, food utensils or preparation surfaces.

#### Re-entry period

**Pre-construction:** Re-entry - **DO NOT** allow entry into uncovered treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after each days use.

**Post-Construction and urban pest control:** Re-entry **DO NOT** allow people and pets to enter treated areas until the spray has dried (normally 3-4 hours) and ventilate buildings before reoccupying. When prior entry is necessary, wear cotton overalls buttoned to the neck, wrist and elbow-length PVC, neoprene or nitrile gloves and chemical resistant footwear. Clothing must be laundered after each days use.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

**Dangerous** to fish and aquatic organisms. **DO NOT** contaminate dams, rivers, streams or waterways or drains with product or used containers.

#### PROTECTION OF PETS AND LIVESTOCK


Before spraying, remove animals and pets from the areas to be treated. Cover or remove any open food and water containers. Cover or remove fish ponds, aquariums etc before spraying.

#### STORAGE, SPILLAGE AND DISPOSAL

Store in closed original container, in a cool, well-ventilated area away from children, animals, food and feedstuffs. **DO NOT** store for prolonged periods in direct sunlight. In case of spillage, confine and absorb spilled product with absorbent material such as sand, clay or cat litter. Dispose of waste as indicated below or according to the Australian Standard AS 2507 - Storage and Handling of Pesticides. **DO NOT** allow spilled product to enter sewers, drains, creeks or any other waterways.

**For REFILLABLE containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**For 1, 5, 10, 20 & 200 L containers:** Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt. **DO NOT** bury waste or surplus product. Dispose of undiluted waste by either dilution and use according to the Directions for Use or returning to the point of purchase in the original container for controlled disposal. Dispose of diluted surplus product by using according to the Directions for Use. **DO NOT** re-use empty container.



**DIRECTIONS FOR USE:**

**RESTRAINTS:**

**DO NOT** use this product at less than indicated label rates.  
**DO NOT** apply to soils if excessively wet or immediately after heavy rain to avoid run-off of the chemical.  
**DO NOT** use in cavity walls (except via certified cavity infill reticulation systems or direct treatment of nest).

PEST	RATE	SITUATIONS	STATE	CRITICAL COMMENTS
Spiders	25-50mL/10L	Internal & external areas & surrounds of domestic, commercial, public & industrial buildings and structures	All States	Refer to Table A
Papernest Wasps	50mL/10L			
Ants (excluding Red imported Fire Ants), cockroaches, mosquitoes, fleas, flies, ticks (excluding the paralysis tick Ixodes holocyclus) (Adults & Nymphs)	50 -100mL/10L			
Subterranean Termites	Refer to Table B	Domestic, Public, Commercial & Industrial areas	All states (except Tas)	Refer to Table C

**TABLE A: CRITICAL COMMENTS for the management and control of a range of interior and exterior urban pests**

<b>Spiders</b>	Use the higher rate in situation where pest pressure is high, when rapid knockdown and/or maximum residual protection is desired. Pay particular attention to protected dark areas such as cracks and crevices, under floors, eaves and other known hiding or resting places. For overall band surface spray, apply as a coarse, low pressure surface spray to areas where spiders hide, frequent and rest. Spray to the point of run-off using around 5L of spray mixtures per 100 m <sup>2</sup> and ensuring thorough coverage of the treated surfaces. For crack & crevice treatment use an appropriate solid stream nozzle. For maximum spider control use a two part treatment. 1. Crack & Crevice 2. Overall band spray surfaces.
<b>Papernest Wasp</b>	Apply prepared solution to the point of run-off directly to the papernest ensuring thorough and even coverage. When all adult wasps have been knocked-down the nest may be safely removed from the structure.
<b>Ants (excluding Red imported Fire Ants), cockroaches, mosquitoes, fleas, flies, ticks (excluding the paralysis tick Ixodes holocyclus) (Adults &amp; Nymphs)</b>	Use the higher rate in situations where pest pressure is high, when rapid knockdown and/or maximum residual protection is desired. The lower rate may be used for follow-up treatments. <b>For indoor use</b> pay particular attention to protected dark areas such as cracks & crevices, behind or under sinks, stoves and refrigerators, furniture, pipes, comices, skirting boards & other known hiding or resting places. <b>DO NOT</b> use as a surface spray. <b>On non-porous surfaces</b> apply as a coarse spray at the rate of 1L solution per 20 m <sup>2</sup> . When treating non-porous surfaces do not exceed the point of run-off. <b>On porous surfaces</b> or use through power equipment, spray at the rate of 1L of solution per 10 m <sup>2</sup> . When treating porous surfaces <b>DO NOT</b> exceed the point of run-off. <b>Ants:</b> To control ants apply to trails and nests. Repeat as necessary. <b>Fleas &amp; Ticks:</b> To control fleas and ticks apply prepared solution to outside surfaces of buildings and surrounds including but not limited to foundations, verandahs, window frames, eaves, patios, garages, pet housing, soil, turf, trunks or woody ornaments or other areas where pests congregate or have been seen. <b>Flies &amp; Mosquitoes:</b> To control flies and mosquitoes apply prepared solution to surfaces where insects rest or harbour. Reapply as necessary. <b>Perimeter treatments:</b> Apply the prepared solution to a band of soil or vegetation two to three metres wide around and adjacent to the structure. Also treat the foundation of the structure to a height of approximately one metre. Use a spray volume of 5 to 10L per 100 m <sup>2</sup> . Higher volumes of water may be needed if organic matter is present or foliage is dense.

**TABLE B: BiFORCE 100SC Termiticide and Insecticide use rates for management of SUBTERRANEAN TERMITES**

SITUATION	All areas SOUTH of the Tropic of Capricorn (except Tas)		All areas NORTH of the Tropic of Capricorn	
	RATE	Expected Protection Period*	RATE	Expected Protection Period*
<b>Pre-Construction Barriers</b> Under slabs and under suspended floors with less than 400 mm crawl space	1L/100L	At least 10 years	1.5L/100L	5 years
	500mL/100L	10 years	1L/100L (Note1)	4 years
			750mL/100L (Note1)	3 years
<b>Perimeter Barriers</b> For new and existing buildings	1L/100L	At least 10 years	1.5L/100L	5 years
	500mL/100L	10 years	1L/100L	4 years
			750mL/100L	3 years
250mL/100L	3 years	500mL/100L	2 years	
<b>Post-Construction Barriers</b> Under slabs and under suspended floors with less than 400 mm crawl space	1L/100L	At least 10 years	1.5L/100L	5 years
	500mL/100L	10 years	1L/100L	4 years
			750mL/100L	3 years
500mL/100L	10 years	500mL/100L	2 years	
<b>Reticulation Systems</b> Perimeter and/or service penetration treatment only	1L/100L	At least 10 years	1.5L/100L	5 years
	500mL/100L	10 years	1L/100L	4 years
			750mL/100L	3 years
250mL/100L	3 years	500mL/100L	2 years	
<b>Reticulation Systems</b> Cavity infill & footing barriers	500mL/100L	5 years	1.25L/100L	2 years
<b>Protection of Poles &amp; Fence Posts</b>	500mL/100L	10 years	1.5L/100L	5 years
			1L/100L	4 years
			750mL/100L	3 years
<b>Nest Eradication</b>	500mL/100L	Not applicable	500mL/100L	Not applicable

\* Several factors contribute to the estimated length of protection provided for each termite treatment. The actual protection period will depend on the termite hazard, climate, soil conditions and rate of termiticide used. The need for retreatment is to be determined as a result of at least an annual inspection, or more frequently in high risk area, by a qualified licensed Pest Control Operator.  
**Note 1:** This rate must be used in conjunction with a certified reticulation system that is capable of distributing the Termiticide solution according to the product label and the Australian Standard AS 3660 Series.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**TABLE C: CRITICAL COMMENTS for use against SUBTERRANEAN TERMITES**

<b>Pre-Construction Barriers</b> Under slabs for protection of new buildings	<ul style="list-style-type: none"> <li>Apply with suitable application equipment to form a complete and continuous chemical barrier (both vertical and horizontal) under the slab. The formation of the barrier may require a combination of conventional open wand application and soil trenching and/or rodding applications. Recommended rod spacing should be between 150 and 300mm, as per soil type. For additional information refer to "CRITICAL APPLICATION DETAILS" on this label and the Australian Standard AS 3660 Series.</li> <li>An external perimeter barrier (both horizontal &amp; vertical) is an essential part of termite protection and must be installed at the completion of the building. Refer to "Perimeter Barriers" below for further details.</li> <li>Chemical barriers that have been disturbed by construction, excavation and/or landscaping activities will need to be reapplied to restore continuity of the barrier.</li> </ul>
<b>Pre-Construction Barriers</b> Under suspended floors	<ul style="list-style-type: none"> <li>For areas under suspended floors with restricted access (typically with less than 400mm clearance), the entire subfloor area should be treated as a continuous horizontal barrier, which completely abuts an internal vertical barrier (if necessary) around any substructure wall. Ideally, this operation should be done during construction of the building while access is more readily available.</li> <li>For areas beneath suspended floors which have adequate access (eg. more than 400 mm clearance), install perimeter barriers around each individual pier, stump, service penetration and substructure wall.</li> <li>An external perimeter barrier (both horizontal &amp; vertical) is an essential part of termite protection and must be installed at the completion of the building. Refer to "Perimeter Barriers" in this leaflet for further details.</li> <li>Chemical barriers that have been disturbed by construction, excavation and/or landscaping activities will need to be reapplied to restore continuity of the barrier.</li> </ul>
<b>Perimeter Barriers</b> For new and existing buildings	<ul style="list-style-type: none"> <li>Perimeter barriers (both horizontal and vertical, external and where required, internal and sub-floor) are an essential part of termite protection and must be installed at the completion of the building. Perimeter barriers should be installed around slabs, piers, substructure walls and external penetrations points.</li> <li>Apply with suitable application equipment to form a continuous chemical barrier (both vertical and horizontal) around the structure and to a depth reaching 80mm below the top of the footings, where appropriate. The formation of the barrier may require a combination of several application techniques, including soil trenching and/or rodding and open wand applications.</li> <li>Chemical barriers that have been disturbed by construction, excavation and/or landscaping activities will need to be reapplied to restore continuity of the barrier.</li> </ul>
<b>Post-Construction Barrier Treatments</b> For the protection of existing buildings	<ul style="list-style-type: none"> <li>Apply with suitable application equipment to form a complete and continuous barrier (both horizontal &amp; vertical) around and under the buildings and structures as in accordance with AS3660 with particular emphasis on any known infestations areas. To form the chemical barrier a number of application techniques may be needed including soil rodding; trenching; open wand and sub-slab injections.</li> <li>Chemical barriers beneath concrete will require drilling. Recommended drill hole spacing is between 150mm and 300 mm. To enhance chemical distribution, use a lateral dispersion tip on the injector and deliver up to 10 L of emulsion per linear meter. Drill holes should be no more than 150 mm from foundation walls or expansion joints to ensure complete formation of a chemical barrier.</li> <li>For areas under suspended floors with restricted access (typically with less than 400 mm clearance), the entire subfloor area should be treated as a continuous horizontal barrier, which completely abuts an internal vertical barrier (if necessary) around any substructure wall. Otherwise, install perimeter barriers around each individual pier, stump, penetration point and structure wall.</li> <li>Chemical barriers that have been disturbed by construction, excavation and/or landscaping activities will need to be reapplied to restore continuity of the barrier.</li> </ul>
<b>Reticulation Systems</b> Perimeter and/or service penetration treatment only	<ul style="list-style-type: none"> <li>BiFORCE 100SC Termiticide &amp; Insecticide must be used through a certified reticulation system to form and replenish perimeter barriers around buildings and service penetrations. The system must be installed according to the manufacturer's specifications and be capable of distributing the termiticide solution according to the product label and the Australian Standard AS 3660 Series.</li> <li>Perimeter barriers consist of a horizontal barrier abutting a vertical barrier, which must reach down to the top of the footing.</li> <li>Delivery pipes must be placed in such a position to ensure that the requirements for both horizontal and vertical barriers as specified in the Australian Standard AS 3660 Series are met. Special attention must also be afforded to the positioning of the delivery pipes to ensure that the resultant termiticidal barriers are continuous and complete.</li> <li>Apply the prepared termiticide solution by pumping through the system according to the manufacturer's specifications. Use a minimum delivery volume of 100L of solution per m<sup>3</sup> of soil. This equates to a delivery volume of 5L of solution per linear metre for a vertical barrier 300 mm x 150 mm in dimension.</li> <li>Pre-Construction - For use in conjunction with full soil treatment horizontal barriers only: Apply the diluted solution through the perimeter reticulation system as specified above. Follow instructions for Pre-construction horizontal barrier formation.</li> </ul>
<b>Reticulation Systems</b> Cavity infill & footing barriers	<ul style="list-style-type: none"> <li>BiFORCE 100SC Termiticide &amp; Insecticide must be used through a certified reticulation system to form and replenish cavity infill and footing barriers. The system must be installed according to the manufacturer's specifications and be capable of distributing the termiticide solution according to the product label and the Australian Standard AS 3660 Series.</li> <li>Delivery pipes must be placed in such a position to ensure that the requirements for both horizontal and vertical barriers as specified in the Australian Standard AS 3660 Series are met. Special attention must also be afforded to the positioning of the delivery pipes to ensure that the resultant termiticidal barriers are continuous and complete.</li> <li>Apply the prepared termiticide solution by pumping through the system according to the manufacturer's specifications with delivery volume of 2L of solution per linear metre of delivery pipe.</li> <li><b>Note: where this system is to be installed at the pre-construction stage, a full under slab pre-construction barrier, applied by either open wand application or suitably certified reticulation system, is also recommended.</b></li> <li>The recommended rate of application is 2L of solution per linear metre which equates to 2L of solution per 0.0068 m<sup>3</sup> or approximately 7L of sand. Should the volume of fill in the wall cavity deviate from 7L (0.17 m x 0.04 m x 1m = 0.0068 m<sup>3</sup>) per linear metre of wall cavity, then the amount of BiFORCE 100SC solution applied per linear metre of wall cavity should be adjusted accordingly. As a guide, the target bifenthrin loading of treated sand/soil in a cavity infill situation is 110 mg/kg South of the Tropic of Capricorn and 220 mg/kg North of the Tropic of Capricorn.</li> <li>To facilitate more even distribution of BiFORCE 100SC solution in the wall cavity, ensure that the fill is evenly compacted at the time of installation. To further enhance distribution saturation of the sand/soil in the infill is recommended at the time of treatment.</li> </ul>
<b>Protection of Service Poles &amp; Fence Posts</b>	<ul style="list-style-type: none"> <li>Create a continuous termiticide barrier 450mm deep and 150mm wide around the pole or post by soil injection or rodding. For new poles and posts, treat backfill and the bottom of the hole. Use 100L of solution per m<sup>3</sup> of soil.</li> <li>Regular inspections should be undertaken to determine when and if treatment is necessary. If disturbance of the barrier has occurred, retreatment of the area affected will be required.</li> <li>Posts and poles may also be drilled and injected with spray solution.</li> <li><b>Note: For existing poles and posts, it is impractical to treat the full depth and underneath of such poles and posts and therefore the possibility of future termite attack from below the treated area cannot be ruled out.</b></li> </ul>
<b>Eradication of Termite Nest</b>	<ul style="list-style-type: none"> <li>Locate nest and flood with insecticide solution. Trees, poles, posts and stumps containing nests may require drilling prior to treatment with termiticide solution. The purpose of drilling is to ensure the termiticide solution is distributed throughout the entire nest. Drill holes in live trees should be sealed with an appropriate caulking compound after injection.</li> </ul>

**NOTE:** The termiticide barrier provided by this product has a finite life. This together with the recommendation to undertake annual inspection must be stated on the durable notice required by the BCA, B1.3 (j) (ii).