

DIRECTIONS FOR USE
DO NOT exceed 3m between bait stations for mice, or 9m for rats. DO NOT place bait stations in the open.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.	Rats (<i>Rattus norvegicus</i> , <i>Rattus rattus</i>)	2 to 8 x 28 g blocks per bait station	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.
	Mice (<i>Mus musculus</i>)	1 to 2 x 28 g blocks per infested area	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

PRECAUTION

The use of this product for the control of protected native rodents requires permission from wildlife authorities.
DO NOT allow bait to contaminate foodstuffs or food intended for human or animal consumption.
DO NOT apply bait to crops.
DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food.
The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

6140-3

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Contrac® Blox®

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

FOR THE CONTROL OF RATS AND MICE

including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

**PRIMARY PACK
CONTAINS 64 x 28 g RODENT BLOCKS**

**BLOCKS ARE ILLEGAL TO SELL SEPARATELY.
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS.**

NET 1.8 kg

Manufactured by:

Bell Laboratories, Inc.
Madison, Wisconsin 53704 U.S.A. • www.belllabs.com

Australian Agent:
Bell Australia Pty Ltd.
Level 4 Grant Thornton House
102 Adelaide Street
Brisbane QLD 4000
Australia

EMERGENCY CONTACT NO. (02) 9647 2111

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

STORAGE AND DISPOSAL

Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in locked room or place away from children, animals, food and foodstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush, puncture and bury empty containers in a local authority landfill. If not available, puncture and bury empty containers in a local specially marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned. Dispose of carcasses safely by burying and burning in an approved landfill.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear elbow-length pvc gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or the Poisons Information Centre (phone Australia 131126; New Zealand 0800 764 766)

MSDS

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE

The use of Contrac® Blox® is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974. 013008

APVMA Approval No.: 48372/1.8KG/0108

Batch No.:

D.O.M.



48372_43806_1.8KG_LABEL_MPL_V2

Total Area: 26.67 cm x 13.2039 cm
100% actual size on A3 dieline

Silk Screen 18#, 4 gallon or 8 kg pail

THIS LINE CAN BE MOVED. CENTER FRONT TO CENTER BACK IS 16"

Handle lug on Finished Pail

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ContraC® Blox®

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

FOR THE CONTROL OF RATS AND MICE

PRIMARY PACK
CONTAINS 285 x 28 g RODENT BLOCKS

BLOCKS ARE ILLEGAL TO SELL SEPARATELY.
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS

Australian Agent:
Bell Australia Pty Ltd.
Level 4 Grant Thornton House
102 Adelaide Street
Brisbane QLD 4000
Australia

EMERGENCY CONTACT NO. (02) 9647 2111

Manufactured by:
Bell Laboratories, Inc.
Madison, Wisconsin 53704 U.S.A. • www.belllabs.com

NET 8 kg



DIRECTIONS FOR USE

DO NOT exceed 3m between bait stations for mice, or 9m for rats. DO NOT place bait stations in the open.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.	Rats (<i>Rattus norvegicus</i> , <i>Rattus rattus</i>) Mice (<i>Mus musculus</i>)	2 to 8 x 28 g blocks per bait station 1 to 2 x 28 g blocks per infested area	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

PRECAUTION

The use of this product for the control of protected native rodents requires permission from wildlife authorities. DO NOT allow bait to contaminate foodstuffs or food intended for human or animal consumption. DO NOT apply bait to crops. DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

ContraC® Blox®

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

FOR THE CONTROL OF RATS AND MICE

including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

PRIMARY PACK
CONTAINS 285 x 28 g RODENT BLOCKS

BLOCKS ARE ILLEGAL TO SELL SEPARATELY.
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS.

NET 8 kg

Manufactured by:
Bell Laboratories, Inc.
Madison, Wisconsin 53704 U.S.A. • www.belllabs.com

Australian Agent:
Bell Australia Pty Ltd.
Level 4 Grant Thornton House
102 Adelaide Street
Brisbane QLD 4000
Australia

EMERGENCY CONTACT NO. (02) 9647 2111

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT
DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

STORAGE AND DISPOSAL
Store in the closed original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in locked room or place away from children, animals, food and feedstuffs, seed and fertilizers. Do not dispose of chemicals on site. Break, crush, puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specially marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Dispose of carcasses safely by burying and burning in an approved landfill.

SAFETY DIRECTIONS
Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID
If poisoning occurs, contact a doctor or the Poisons Information Centre (phone Australia 131126; New Zealand 0800 764 766)

MSDS
For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE
The use of ContraC® Blox® is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974. 050308

APVMA Approval No.: 48372/BKG/0508

Batch No.:

D.O.M.

48372_43806_8KG_LABEL_MPL_V4

Total Area: 69.84175 cm x 19.4667 cm
58% to 100% actual size on A3 dieline

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONTRAC[®] BLOX[™]



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)


For the control of rats and mice, including those resistant to other anticoagulants,
in and around domestic, commercial, public services, and industrial buildings,

PRIMARY PACK

CONTAINS 20 x 450 G RODENT BLOCKS WHICH ARE ILLEGAL TO SELL
SEPARATELY.
DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS.

NET 9 kg

<i>Manufactured by:</i> BELL LABORATORIES, INC. MADISON, WI 53704 U.S.A. www.belllabs.com	<i>Australian Agent:</i> Bell Australia Pty Ltd. Level 4 Grant Thornton House 102 Adelaide Street Brisbane QLD 4000 AUSTRALIA PH: 02 9647 2111
--	--



DIRECTIONS FOR USE

DO NOT exceed 3m between bait stations for mice, or 9m for rats. DO NOT place in the open.

Situation	Pest	Rate	Critical Comments
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships), and around grain terminals.	Rats (<i>Rattus rattus</i> , <i>Rattus norvegicus</i>)	1 block per bait station	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.
	Mice (<i>mus musculus</i>)	1 block per infested area	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.


PRECAUTION

The use of this product for the control of protected native rodents requires permission from wildlife authorities.

DO NOT allow bait to contaminate foodstuffs or food intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food. The product is restricted to use in and around buildings (within 2 metres), or enclosed spaces.



PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

STORAGE AND DISPOSAL

Store in the closed original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in locked room or place away from children, animals, food and feedstuffs, seed and fertilizers. Do not dispose of chemicals on site. Break, crush, puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specially marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt. Dispose of carcasses safely by burying and burning in an approved landfill.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear elbow-length pvc gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K₁ (Phytomenadione) is antidotal.

FIRST AID

If poisoning occurs, contact a doctor or the Poisons Information Centre (phone Australia 131126; New Zealand 0800 764 766)

MSDS

For further information please refer to the Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE:

The use of CONTRAC® Blox® is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories Inc., regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save those nonexcludable conditions implied by the Trade Practices Act, 1974.

APVMA Approval No.: 48372/9KG/0108

Batch No.:
D.O.M.

Contrac is a trademark of Bell Laboratories, Inc., U.S.A.

